




Drug	Dose and Administration	Renal Dose Adjustments	Key Drug–Drug Interactions
Nirmatrelvir + ritonavir	Adults and children aged ≥12 years: 300 mg/100 mg PO twice daily x 5 days	If eGFR ≥30 to <60 mL/min, use 150 mg/100 mg PO twice daily x 5 days If eGFR <30 mL/min, do not use	Common: CYP3A
Remdesivir	Adults and children (weight ≥40 kg): 200 mg IV loading dose on Day 1, then 100 mg on Days 2 and 3 Children (aged ≥28 days and weight ≥3 kg to <40 kg): 5 mg/kg IV loading dose on Day 1, then 2.5 mg/kg on Days 2 and 3	If eGFR <30 mL/min, use is not recommended	Chloroquine phosphate and hydroxychloroquine sulfate
Molnupiravir	Adults (aged ≥18 years): 800 mg PO every 12 hours x 5 days	None	None

CYP, cytochrome P450; eGFR, estimated glomerular filtration rate.

References

covid19treatmentguidelines.nih.gov/management/clinical-management-of-adults/nonhospitalized-adults--therapeutic-management.
 covid19treatmentguidelines.nih.gov/management/clinical-management-of-children/nonhospitalized-children-therapeutic-management.
 Molnupiravir PI.
 Nirmatrelvir/ritonavir PI.
 Remdesivir PI.